

Please write clearly in block capitals.	
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	

# AS **MATHEMATICS**

Paper 1

Wednesday 15 May 2019

Morning

Time allowed: 1 hour 30 minutes

### **Materials**

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

## Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer each question in the space provided for that question.
   If you require extra space, use an AQA supplementary answer book; do not use the space provided for a different question.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

# Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

# Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Exam	iner's Use
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
TOTAL	
· · · · · · · · · · · · · · · · · · ·	



# Section A

Answer all questions in the spaces provided.

State the number of solutions to the equation  $\tan 4\theta = 1$  for  $0^{\circ} < \theta < 180^{\circ}$ 1 Circle your answer.

[1 mark]

1

2

8

Dan believes that 2

for every positive integer n, at least one of  $2^n - 1$  and  $2^n + 1$  is prime.

Which value of n shown below is a counter example to Dan's belief?

Circle your answer.

[1 mark]

$$n = 3$$

$$n = 2$$

$$n=4$$
  $n=5$ 

$$n = 6$$

3	It is given that $(x + 1)$ and $(x - 3)$ are two factors of $f(x)$ , where	
	$f(x) = px^3 - 3x^2 - 8x + q$	
3 (a)	Find the values of $p$ and $q$ .	[3 marks]
3 (b)	Fully factorise f (x).	[2 marks]



4	Show that $\frac{\sqrt{6}}{\sqrt{3}-\sqrt{2}}$ can be expressed in the form $m\sqrt{n}+n\sqrt{m}$ , where $m$ a integers.	and <i>n</i> are
	Fully justify your answer.	[4 marks]



5	(a)	Sketch	the	CUIVE	v = g(x)	where
J 1	(a)	OKELLII	เมเษ	cuive	$V - \mathbf{g}(\lambda)$	WITCIE

$$g(x) = (x + 2)(x - 1)^2$$

[3 marks]

<b>5 (b)</b> Her	nce, solve $g(x) \leq 0$		[2 marks]

0 5

Turn over ▶

6 (a) (i)	Show that $\cos \theta = \frac{1}{2}$ is one solution of the equation	
	$6\sin^2\theta + 5\cos\theta = 7$	[2 marks]
6 (a) (ii)	Find all the values of $\theta$ that solve the equation	
	$6\sin^2\theta + 5\cos\theta = 7$	
	for $0^{\circ} \le \theta \le 360^{\circ}$	
	Give your answers to the nearest degree.	[2 marks]



6 (b)	Hence, find all the solutions of the equation	
	$6\sin^2 2\theta + 5\cos 2\theta = 7$	
	for $0^{\circ} \leq \theta \leq 360^{\circ}$	
	Give your answers to the nearest degree.	[2 marks]

Turn over for the next question

7	Given that $y \in \mathbb{R}$ , prove that	
	$(2+3y)^4+(2-3y)^4\geq 32$	
	Fully justify your answer.	[6 marks]



8	Prove that the curve with equation					
	$y = 2x^5 + 5x^4 + 10x^3 - 8$					
	has <b>only one</b> stationary point, stating its coordinates.	[6 marks]				
	Turn over for the next question					



, ,	curve cuts the $x$ -axis at $(2, 0)$	anu n	as gradier	it function		
		$\frac{\mathrm{d}y}{\mathrm{d}x} =$	$=\frac{24}{x^3}$			
Fi	ind the equation of the curve.					[4 m
_						
_					<del></del>	
_						
_						
_						
_						
_						
_						
_					····	
_						



normal to the curve at (2, 0)		



On 18 March 2019 there were 12 hours of daylight in Inverness.		
On 16 June 2019, 90 days later, there will be 18 hours of daylight in Invernes	S.	
Jude decides to model the number of hours of daylight in Inverness, $N,$ by the formula	е	
$N = A + B \sin t^{\circ}$		
where $t$ is the number of days after 18 March 2019.		
State the value that Jude should use for $A$ .	[1 r	mark]
State the value that Jude should use for <i>B</i> .	[1 ɪ	mark]
Using Jude's model, calculate the number of hours of daylight in Inverness or		
15 May 2019, 58 days after 18 March 2019.		mark]
	On 16 June 2019, 90 days later, there will be 18 hours of daylight in Inverness Jude decides to model the number of hours of daylight in Inverness, $N$ , by the formula $N = A + B \sin t^\circ$ where $t$ is the number of days after 18 March 2019. State the value that Jude should use for $A$ .  State the value that Jude should use for $B$ .	On 16 June 2019, 90 days later, there will be 18 hours of daylight in Inverness. Jude decides to model the number of hours of daylight in Inverness, $N$ , by the formula $N = A + B \sin t^\circ$ where $t$ is the number of days after 18 March 2019. State the value that Jude should use for $A$ . [1 Inverse of the value that Jude should use for $B$ . [1 Inverse of the value that Jude should use for $B$ .]

10 (a) (iv)	Using Jude's model, find how many days during 2019 will have at least 17.4 hours of daylight in Inverness.		
	[4 marks]		
10 (a) (v)	Explain why Jude's model will become inaccurate for 2020 and future years.  [1 mark]		
10 (b)	Anisa decides to model the number of hours of daylight in Inverness with the formula		
	$N = A + B \sin\left(\frac{360}{365}t\right)^{\circ}$		
	Explain why Anisa's model is better than Jude's model.  [1 mark]		



	Section B
	Answer <b>all</b> questions in the spaces provided.
11	A ball moves in a straight line and passes through two fixed points, A and B, which are 0.5 m apart.
	The ball is moving with a constant acceleration of $0.39\mathrm{ms^{-2}}$ in the direction AB.
	The speed of the ball at $A$ is $1.9\mathrm{ms^{-1}}$
	Find the speed of the ball at B.
	Circle your answer.  [1 mark]
	$2 \mathrm{ms^{-1}}$ $3.2 \mathrm{ms^{-1}}$ $3.8 \mathrm{ms^{-1}}$ $4 \mathrm{ms^{-1}}$
	21115 3.21115 3.01115 41115
12	A particle $P$ , of mass $m$ kilograms, is attached to one end of a light inextensible string.
	The other end of this string is held at a fixed position, O.
	P hangs freely, in equilibrium, vertically below O.
	Identify the statement below that correctly describes the tension, $T$ newtons, in the string as $m$ varies.
	Tick (✓) one box.
	[1 mark]
	T varies along the string, with its greatest value at O
	T varies along the string, with its greatest value at P
	$T={ m 0}$ because the system is in equilibrium
	T is directly proportional to m



Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

1 5

13	A car, starting from rest, is driven along a horizontal track.	
	The velocity of the car, $v  \text{m}  \text{s}^{-1}$ , at time $t$ seconds, is modelled by the equa	tion
	$v = 0.48t^2 - 0.024t^3$ for $0 \le t \le 15$	
13 (a)	Find the distance the car travels during the first 10 seconds of its journey.	[3 marks]
13 (b)	Find the maximum speed of the car.	
10 (5)	Give your answer to three significant figures.	[4 marks]
		[



13 (c)	Deduce the range of values of $t$ for which the car is modelled as decelerating. [2 marks]
	Turn over for the next question



14	Two particles, $A$ and $B$ , lie at rest on a smooth horizontal plane.	
	A has position vector $\mathbf{r}_A = (13\mathbf{i} - 22\mathbf{j})$ metres B has position vector $\mathbf{r}_B = (3\mathbf{i} + 2\mathbf{j})$ metres	
14 (a)	Calculate the distance between A and B.	[2 marks]
14 (b)	Three forces, $F_1$ , $F_2$ and $F_3$ are applied to particle $A$ , where	
()	$\mathbf{F_1} = (-2\mathbf{i} + 4\mathbf{j})$ newtons	
	$\mathbf{F}_2 = (6\mathbf{i} - 10\mathbf{j}) \text{ newtons}$	
	Given that $A$ remains at rest, explain why $\mathbf{F}_3 = (-4\mathbf{i} + 6\mathbf{j})$ newtons	
		[1 mark]



14 (c)	A force of $(5\mathbf{i} - 12\mathbf{j})$ newtons, is applied to $B$ , so that $B$ moves, from rest, in a straig line towards $A$ .	ght
	B has a mass of 0.8 kg	
14 (c) (i)	Show that the acceleration of $B$ towards $A$ is $16.25\mathrm{ms^{-2}}$	ks]
		_
		_
		_
		_
14 (c) (ii)	Hence, find the time taken for <i>B</i> to reach <i>A</i> .	
	Give your answer to two significant figures.  [2 mar	ks]
		_
		_
		_
		-




A tractor and its driver have a combined mass of $m$ kilograms.
The tractor is towing a trailer of mass $4m$ kilograms in a straight line along a horizontal road.
The tractor and trailer are connected by a horizontal tow bar, modelled as a light rigid rod.
A driving force of 11 080 N and a total resistance force of 160 N act on the tractor.
A total resistance force of 600 N acts on the trailer.
The tractor and the trailer have an acceleration of $0.8\mathrm{ms^{-2}}$
Find m. [3 marks]
Find the tension in the tow bar.  [2 marks]




15 (c)	At the instant the speed of the tractor reaches $18\mathrm{km}\mathrm{h}^{-1}$ the tow bar breaks.		
	The total resistance force acting on the trailer remains constant.		
	Starting from the instant the tow bar breaks, calculate the time taken until the speed of the trailer reduces to $9\mathrm{km}h^{-1}$		
	[4 marks]		
	END OF QUESTIONS		
	2.12 0. 4020110110		











There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED Copyright information For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Do not write outside the box

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.



