Time allowed: 1 hour 45 minutes



GCSE CHEMISTRY



Higher Tier Chemistry 1H

Specimen 2018

Materials

For this paper you must have:

- a ruler
- a calculator
- the periodic table (enclosed).

Instructions

- Answer all questions in the spaces provided.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- There are 100 marks available on this paper.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.
- When answering questions 02.3, 05.2, 08.5 and 09.4 you need to make sure that your answer:
 - is clear, logical, sensibly structured
 - fully meets the requirements of the question
 - shows that each separate point or step supports the overall answer.

Advice

In all calculations, show clearly how you work out your answer.

Please write clearly, in block capitals.					
Centre number Candidate number Candidate number					
Surname					
Forename(s)					
Candidate signature					

This question is about halogens and their compounds.

Table 1 shows the boiling points and properties of some of the elements in Group 7 of the periodic table.

Table 1

Element	Boiling point in [°] C	Colour in aqueous solution
Fluorine	-188	colourless
Chlorine	-35	pale green
Bromine	Х	orange
Iodine	184	brown

0 1 . 1	Why does iodine have a higher boiling point than chlorine?			
	Tick one box.		[1 mark]	
	lodine is ionic and chlorine is covalent			
	lodine is less reactive than chlorine			
	The covalent bonds between iodine atoms are stronger			
	The forces between iodine molecules are stronger			
0 1 . 2	Predict the boiling point of bromine.		[1 mark]	

A redox reaction takes place when aqueous chlorine is added to potassium iodide solution.

The equation for this reaction is:

$$Cl_2(aq) + 2KI(aq) \rightarrow I_2(aq) + 2KCI(aq)$$

0 1 . 3	Look at Table 1 .		
	What is the colour of the final solution	in this reaction?	[4 manula]
	Tick one box.		[1 mark]
	Brown		
	Orange		
	Pale green		
	Colourless		
0 1 . 4	What is the ionic equation for the read	ction of chlorine with potassium iodide	?
	Tick one box.	·	[1 mark]
	$Cl_2 + 2K \rightarrow 2KCI$		
	$2l^- + Cl_2 \rightarrow l_2 + 2Cl^-$		
	$I^- + CI \rightarrow I + CI^-$		
	$I^- + K^+ \rightarrow KI$		

Question 1 continues on the next page

Why does potassium iodide solution conduct electricity?					
Tick one box.					
It contains a metal					
It contains electrons which can move					
It contains ions which can move					
It contains water					
What are the products of electric one box.	ctrolysing potassium iodide solution	n?	[1 mark]		
Product at cathode	Product at anode				
Product at cathode hydrogen	Product at anode iodine				
hydrogen	iodine				
	Tick one box. It contains a metal It contains electrons which can medit contains ions which can medit contains water What are the products of electrons	Tick one box. It contains a metal It contains electrons which can move It contains ions which can move It contains water What are the products of electrolysing potassium iodide solution	Tick one box. It contains a metal		

0 2	An atom of aluminium has the symbol ²⁷ ₁₃ Al						
0 2 . 1	Give the number of protons, neutrons and electrons in this atom of aluminium						
		[3 marks]					
	Number of protons Number of neutrons Number of electrons						
0 2 . 2	Why is aluminium positioned in Group 3 of the periodic table?						
		[1 mark]					

Question 2 continues on the next page

0 2 . 3 In the periodic table, the transition elements and Group 1 elements are metals.

Some of the properties of two transition elements and two Group 1 elements are shown in

Table 2.


Table 2

	Transitio	n elements	Group 1	elements
	Chromium	Iron	Sodium	Caesium
Melting point in °C	1857	1535	98	29
Formula of oxides	CrO Cr_2O_3 CrO_2 CrO_3	FeO Fe₂O₃ Fe₃O₄	Na₂O	Cs₂O

Use your own knowledge and the data in Table 2 to compare the chemical physical properties of transition elements and Group 1 elements.		
projection of transfer comments and Group it comments.	[6 marks]	

O 3 Figure 1 shows the outer electrons in an atom of the Group 1 element potassium and in an atom of the Group 6 element sulfur.

Figure 1



0 3 . 1	Potassium forms an ionic compound with suitur.
	Describe what happens when two atoms of potassium react with one atom of sulfur.

Give your answer in terms of electron transfer.


Give the formulae of the ions formed.

[5 marks]

Question 3 continues on the next page

0 3 . 2 The structure of potassium sulfide can be represented using the ball and stick model in Figure 2.

Figure 2



The ball and stick model is **not** a true representation of the structure of potassium sulfide.

Give one reason why.


11	ma	rk1

n	3	3	Sulfur	can	also	form	covalent	honds
v	J		Sullul	Call	aisu	IUIIII	COVAICIL	DUITUS

Complete the dot and cross diagram to show the covalent bonding in a molecule of hydrogen sulfide.

Show the outer shell electrons only.

[2 marks]



$\boxed{\mathbf{0} \ \mathbf{3}}$. $\boxed{\mathbf{4}}$ Calculate the relative formula mass (M_r) of aluminium sulfate, $\mathrm{Al}_2(\mathrm{SO}_4)_3$

Relative atomic masses (A_r): oxygen = 16; aluminium = 27; sulfur = 32

[2	marks]

Relative formula mass =

Question 3 continues on the next page

0 3 . 5 Covalent compounds such as hydrogen sulfide have low melting points and do **not** conduct electricity when molten.

Draw **one** line from each property to the explanation of the property.

[2 marks]

Property

Low melting point

Does not conduct electricity when molten

Explanation of property

Electrons are free to move

There are no charged particles free to move

lons are free to move

Weak intermolecular forces of attraction

Bonds are weak

Bonds are strong

0 3 . 6 Ionic compounds such as potassium sulfide have high boiling points and conduct electricity when dissolved in water.

Draw **one** line from each property to the explanation of the property.

[2 marks]


Property Explanation of property Electrons are free to move There are no charged particles free to move High boiling point lons are free to move Weak intermolecular forces of attraction Conduct electricity when molten Bonds are weak Bonds are strong

Turn over for the next question

0 4	Rock salt is a mixture of sand and salt.	
	Salt dissolves in water. Sand does not dissolve in water.	
	Some students separated rock salt.	
	 This is the method used. Place the rock salt in a beaker. Add 100 cm³ of cold water. Allow the sand to settle to the bottom of the beaker. Carefully pour the salty water into an evaporating dish. Heat the contents of the evaporating dish with a Bunsen burner until salt crystals start to form. 	
0 4 . 1	Suggest one improvement to step 2 to make sure all the salt is dissolved in the water. [1 main	· k]
0 4 . 2	The salty water in step 4 still contained very small grains of sand. Suggest one improvement to step 4 to remove all the sand. [1 main	·k]
0 4 . 3	Suggest one safety precaution the students should take in step 5. [1 mail]	·k]

Another student removed water from salty water using the apparatus in Figure 3.

Figure 3



0	4] .	4	Describe how this technique works by referring to the processes a	at A and B .
					[2 marks]
0	4	. [5	What is the reading on the thermometer during this process?	
					[1 mark]
					°C

0 5	A student investigated the reactions of copper carbonate and copper oxide with dilute hydrochloric acid.
	In both reactions one of the products is copper chloride.
0 5 . 1	Describe how a sample of copper chloride crystals could be made from copper carbonate and dilute hydrochloric acid. [4 marks]

0 5 . 2	A student wanted to make 11.0 g of copper chloride.	
	The equation for the reaction is:	
	$CuCO_3 + 2HCI \rightarrow CuCl_2 + H_2O + CO_2$	
	Relative atomic masses, A_r : H = 1; C = 12; O = 16; CI = 35.5; Cu	= 63.5
	Calculate the mass of copper carbonate the student should react with di hydrochloric acid to make 11.0 g of copper chloride.	ilute [4 marks]
	Mass of copper carbonate =	g
0 5 . 3	The percentage yield of copper chloride was 79.1 %.	
	Calculate the mass of copper chloride the student actually produced.	[2 marks]
	Actual mass of copper chloride produced =	g

SPECIMEN MATERIAL Turn over ▶

Question 5 continues on the next page


0 5 . 4	Look at the equations for the two reactions:	
	Reaction 1 $CuCO_3(s) + 2HCI(aq) \rightarrow CuCI_2(aq) + H_2O(I) + CO_2(g)$ Reaction 2 $CuO(s) + 2HCI(aq) \rightarrow CuCI_2(aq) + H_2O(I)$	
	Reactive formula masses: $CuO = 79.5$; $HCI = 36.5$; $CuCI2 = 134.5$; $H2O = 18$	
	The percentage atom economy for a reaction is calculated using: Relative formula mass of desired product from equation × 100 Sum of relative formula masses of all reactants from equation	
	Calculate the percentage atom economy for Reaction 2. [3 marks]	
	Percentage atom economy = %	
0 5 . 5	The atom economy for Reaction 1 is 68.45 %. Compare the atom economies of the two reactions for making copper chloride.	
	Give a reason for the difference. [1 mark]	

Turn over for the next question

0 6

A student investigated simple cells using the apparatus shown in Figure 4.

Figure 4



- If metal 2 is more reactive than metal 1 then the voltage measured is positive.
- If metal 1 is more reactive than metal 2 then the voltage measured is negative.
- The bigger the difference in reactivity of the two metals, the larger the voltage produced.

The student's results are shown in Table 3.

Table 3

Metal 2 Metal 1	Chromium	Copper	Iron	Tin	Zinc
Chromium	0.0 V				
Copper	1.2 V	0.0 V			
Iron	0.5 V	not measured	0.0 V		
Tin	0.8 V	-0.4 V	0.3 V	0.0 V	
Zinc	0.2 V	-1.0 V	-0.3 V	-0.6 V	0.0 V

0 6 . 1	The ionic equation for the reaction occuring at the zinc electrode in the simple cell made using copper and zinc electrodes is:		
	$Zn \rightarrow Zn^{2+} + 2e^{-}$		
	Zinc is oxidised in this reaction.		
	Give a reason why this is oxidation. [1 mark]		
0 6 . 2	Look at Table 3 .		
	Which one of the metals used was the least reactive?		
	Give a reason for your answer. [2 marks]		
	Metal		
	Reason		

Question 6 continues on the next page

0 6 . 3	Predict the voltage that would be obtained for a simple cell that has iron as metal 1 and copper as metal 2 .
	Explain your answer. [3 marks]
0 6 . 4	Hydrogen fuel cells have been developed for cars.
	Write a word equation for the overall reaction that takes place in a hydrogen fuel cell. [1 mark]
0 6 . 5	Write the two half equations for the reactions that occur at the electrodes in a hydrogen fuel cell.
	[2 marks]

Turn over for the next question

0 7 Sodium carbonate reacts with dilute hydrochloric acid:


$$Na_2CO_3 + 2HCI \rightarrow 2NaCI + H_2O + CO_2$$

A student investigated the volume of carbon dioxide produced when different masses of sodium carbonate were reacted with dilute hydrochloric acid.

This is the method used.

- 1. Place a known mass of sodium carbonate in a conical flask.
- 2. Measure 10 cm³ of dilute hydrochloric acid using a measuring cylinder.
- 3. Pour the acid into the conical flask.
- 4. Place a bung in the flask and collect the gas until the reaction is complete.
- 0 7 . 1 The student set up the apparatus as shown in Figure 5.

Figure 5



Identify the error in the way the student set up the apparatus.

Describe what would happen if the student used the apparatus shown.

[2 marks]

The student corrected the error.

The student's results are shown in Table 4.

Table 4

Mass of sodium carbonate in g	Volume of carbon dioxide gas in cm ³
0.07	16.0
0.12	27.5
0.23	52.0
0.29	12.5
0.34	77.0
0.54	95.0
0.59	95.0
0.65	95.0

0 7 . 2	The result for 0.29 g of sodium carbonate is anomalous.		
	Suggest what may have happened to cause this anomalous result. [1 m	nark]	
0 7 . 3	Why does the volume of carbon dioxide collected stop increasing at 95.0 cm ³ ? [1 n	nark]	

Question 7 continues on the next page

0 7 . 4	What further work could the student do to be more certain about the minimum mass of sodium carbonate needed to produce 95.0 cm ³ of carbon dioxide?
	[1 mark]
0 7 . 5	The carbon dioxide was collected at room temperature and pressure. The volume of one mole of any gas at room temperature and pressure is 24.0 dm ³ .
	How many moles of carbon dioxide is 95.0 cm ³ ?
	Give your answer in three significant figures. [2 marks]
	mol
0 7 . 6	Suggest one improvement that could be made to the apparatus used that would give more accurate results.
	Give a reason for your answer.
	[2 marks]

0 7 . 7		nt said that the results of the gas collected were air.	he experiment were wrong bec	ause the first few
	A second s	student said this would ma	ke no difference to the results.	
	Explain why	ny the second student was	correct.	[2 marks]

Turn over for the next question

0 8	Sodium hydroxide neutralises sulfuric acid.
	The equation for the reaction is:
	2NaOH + $H_2SO_4 \rightarrow Na_2SO_4 + 2H_2O$
0 8 . 1	Sulfuric acid is a strong acid.
	What is meant by a strong acid? [2 marks]
08.2	Write the ionic equation for this neutralisation reaction. Include state symbols. [2 marks]

A student used a pipette to add 25.0 cm³ of sodium hydroxide of unknown concentration to a conical flask.

The student carried out a titration to find out the volume of 0.100 mol/dm³ sulfuric acid needed to neutralise the sodium hydroxide.

0 8 . 3	Describe how the student would complete the titration.				
	You should name a suitable indicator and give the colour change that would be seen. [4 marks]				

Question 8 continues on the next page

The student carried out five titrations. Her results are shown in **Table 5**.

Table 5

	Titration 1	Titration 2	Titration 3	Titration 4	Titration 5
Volume of 0.100 mol/dm ³ sulfuric acid in cm ³	27.40	28.15	27.05	27.15	27.15

08.4	Concordant results are within 0.10 cm ³ of each other.	
	Use the student's concordant results to work out the mean volume of 0.100 mol/dm ³	3
	sulfuric acid added. [2 mark	s]
	Mean volume = cm ³	}

The equation for the reaction is:	
$2NaOH + H_2SO_4 \rightarrow Na_2SO_4 + 2H_2O$	
Calculate the concentration of the sodium hydroxide.	
Give your answer to three significant figures.	[4 marks]
Concentration =	mol/dm ³
The student did another experiment using 20 cm ³ of sodium hydroxide s a concentration of 0.18 mol/dm ³ .	olution with
Relative formula mass (M_r) of NaOH = 40	
Calculate the mass of sodium hydroxide in 20 cm ³ of this solution.	[2 marks]
Mass =	g
	$2\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$ Calculate the concentration of the sodium hydroxide. Give your answer to three significant figures.

Turn over for the next question

0 9 This question is about the reaction of ethene and bromine.

The equation for the reaction is:

$$C_2H_4 + Br_2 \rightarrow C_2H_4Br_2$$


0 9 . 1 Complete the reaction profile in Figure 6.

Draw labelled arrows to show:

- The energy given out (ΔH)
- The activation energy.

[3 marks]

Figure 6



0 9 . 2	When ethene reacts with bromine, energy is required to break covalent bonds in the
	molecules

Explain how a covalent bond holds two atoms together.

[2 marks]

Figure 7 shows the displayed formulae for the reaction of ethene with bromine.

Figure 7

The bond enthalpies and the overall energy change are shown in **Table 6**.

Table 6

	C=C	C-H	C-C	C-Br	Overall energy change
Energy in kJ/mole	612	412	348	276	-95

	calculate the bond energy for the [3 marks]
Bond energy	kJ/mole

0 9 . 4 Figure 8 shows the reaction between ethene and chlorine and is similar to the reaction between ethene and bromine.

Figure 8

$$H = C + CI - CI \longrightarrow H - C - C - H$$

$$H = C - C - H$$

$$CI = CI$$

"The more energy levels (shells) of electrons an atom has, the weaker the covalent bonds that it forms."

Use the above statement to predict and explain how the overall energy change for the reaction of ethene with chlorine will differ from the overall energy change for the reaction of ethene with bromine.

[6 marks]

END OF QUESTIONS

There are no questions printed on this page

There are no questions printed on this page

Copyright information

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements in future papers if notified. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2016 AQA and its licensors. All rights reserved.